

ValleyOrtho Rehabilitation Playbook Series

Physician: Dr. Ferdinand Liotta
Office Phone: 970-384-7637

Physician Assistant: Amanda Hunter & Erin Flores **MA:** Daisy Herrera **ATC:** Maddy Prior
Office Fax: 970-384-8133

Surgical Procedure: Meniscectomy

The intent of this information is to inform the treating clinician on the evidence-based considerations to be used as a guideline regarding the surgery noted above. This is not a substitute for appropriate clinical decision making, but a supplement to that effect. If at any time a clinician feels uncertain about a given phase discrepancy or patient presentation they are strongly encouraged to discuss this with the referring physician and his/her team.

****It is the responsibility of the therapist to read the operative report before providing care to the patient to improve treatment communication***.*

Therapeutic Activity Progression Disclaimer: Progression to the next phase should be strongly based on meeting clinical criteria (not solely based on the post-operative timeframes) as appropriate and in collaboration with the referring surgeon. Exercise prescription should be clinically directed by pain and performance absent of detrimental movement patterns with respect to proper biomechanics of the spine, hip, knee and ankle.

Communication Recommendations from Therapist to Surgical

Team: When a treating therapist feels the need to reach out to Dr. Liotta, or a member of his team, at any point for any reason they are strongly encouraged to do so. All concerns are not explicitly written and clinical judgement is paramount. Below is a handful of reasons and suggested methods of contact to promote communication:

Urgent Red Flag Communication

- Uncontrollable and unremitting pain.
- Signs of infection at incision or treated limb.
- Severe palpation tenderness, swelling, tachycardia (UE or LE DVT).
- Labored breathing (PE).
- Drastic decline in ROM.
- After a fall/trauma, or near fall/trauma, resulting in a clinical change.

Preferred Contact Method: 1. Immediate call to MD or PA Cell.
2. Office phone call to request consult with MD/PA/MA/ATC until answer.

Administrative Needs

- Rehabilitation Prescription needed or specific Rx change requests.
- Appointment needed with the physician office, or medication refill.

Preferred Contact Method: Office phone call to MA/ATC.

Other Patient Concerns During Clinic Hours M-TH 9am-5pm F 9-3pm

- Abnormal pain, comorbidities or complications that may prevent attainment of established discharge criteria.
- Patient is noncompliant with rehabilitation process.
- Excessive muscle guarding/motion phobia after 1-2 outpatient visits.
- Adverse work or home practices negatively impacting recovery.
- Patient expresses discontent or concerns with the current POC established by PT and/or by MD/PA

Preferred Contact Method: Phone call to MD &/or PA

Preferred Updates before checkup visits with MD/PA

During Clinic Hours M-TH 9am-5pm F 9-3pm

- Information regarding adherence/participation in rehabilitation process.
- Comments on progress and trend of the patient's rehab course.

Preferred Contact Method: Phone call MD and/or PA. Or Fax update.



Phase 1: Edema, Quadriceps & ROM Recovery (wks 0 -2)

Goals:

- Initiate therapy around post-op day 4¹⁰
- Minimize pain/swelling to decrease quad inhibition^{2,9}
- Normalize quadriceps activation/control^{2,9}
- Set baseline KOOS-pain/KOOS-Sport for RTS readiness¹ (Appendix A)

Precautions/Restrictions:

- WB/Gait:
 - WBAT^{2,9,12}, initial ambulation with bilateral crutches^{2,9}
 - Wean patient from crutches ≠ limp and pain as able^{2,9}
- A/AA/PROM:
 - Week 0-1 ROM 0-90^{02,9} Then ROM progression as tolerated for full ROM by wks 4-6⁹
 - Emphasis should be towards terminal knee extension initially^{2,9}
- Activity:
 - No impact training until week 4+^{9,11}

Phase 1 Therapeutic Activities:

- Gait:
 - Progression from bilateral crutches to single crutch to no AD as able
- ROM:
 - Manual & self-management for flexibility, swelling and full ext^{2,9}
 - Scar and patellar mobilizations on healed incisions⁹
 - Bike partial or full revolutions for ROM gains/maintenance^{2,9}
- Strengthening:
 - Total lower extremity OKC and CKC strengthening/activities aimed avoid valgus collapse and promote core strength/pelvis control through full knee ROM as tolerated^{2,9}
 - Quad TKE focused activity^{2,9}
 - NMES to quad with volitional contraction as needed^{2,9}
 - Slow and progressive NWB total LE strengthening²
 - Mini Squats TKE to mid-range to tolerance^{2,9}
- Balance:
 - Proprioception with TKE control^{2,9,13}

Minimum Criteria for Progression to Phase 2:

- AROM 0⁰-90⁰²
- 20 SLR ≠ Quad Lag^{2,14}

Phase 2: ROM & Total LE Strengthening (wks 3 - 4)

Goals:

- Rehabilitation may progress aggressively because there is no anatomic structure that requires protection^{2,3,10} while concurrently controlling for effusion, pain and inflammation³
- Restore near full ROM^{2,9}
- Normalize gait without AD²
- Improve muscle strength and endurance²
- Improve balance and proprioception^{2,9,13}

Precautions:

- Activity:
 - Avoid impact training until weeks 4+^{9,12}
 - Once single leg press is ≥ 75% LSI begin submax impact training with progressions as tolerated to full intensity impact activities⁹

Phase 2 Therapeutic Activities:

- Gait:
 - Ensure proper weight shifting over involved extremity
- ROM:
 - Manual & self-management for flexibility, swelling for return to full ROM⁹
- Strengthening:
 - Total lower extremity OKC and CKC strengthening/activities aimed avoid valgus collapse and promote core strength/pelvis control through full knee ROM as tolerated^{2,9}
 - High-load progressive quad strengthening may be indicated to target an increase in muscle activation specifically in concentric actions at 70-80% 1 repetition max^{7,10}
 - Recovery of hamstring muscle function and quadriceps control at low force production levels during stability training is not as indicated as concentric quad strength recovery⁷
- Balance:
 - Proprioception training progressions^{2,9,13}

Criteria for Progression to Phase 3:

- Normal gait mechanics without AD
- 0-125° AROM



Phase 3: Total LE Strengthening & Return to Activity (wks 5+)

Goals:

- Address remaining barriers to RTS via KOOS-pain/KOOS-sport¹
- Optimize biomechanics at the hip, knee and ankle
- Increasing strength to support desired activity
- **In prepubescent patients:** focus primarily on form control and movement patterns instead of muscle hypertrophy as their bodies will not put on muscle growth as in more mature patients¹⁵
- Establish patient specific HEP relative to resources and goals.

Precautions:

Phase 3 Therapeutic Activities:

- **ROM:**
 - Manual & self-management for gains in ROM, flexibility & swelling
- **Strengthening & Activity; As Tolerated:**
 - Running Progressions with proper swelling and pain control¹¹
 - Slow progressions of cutting/pivot & decelerating intensity⁹
 - Continue total lower extremity strengthening based on deficits
 - Neuromuscular training for proper landing mechanics is important as patient's alter jump landing mechanics to decrease quad use (forward anterior trunk lean with or without increased knee flexion) in involved knee up to 3 months after surgery despite = Quad LSI muscle testing⁶
- **Balance:**
 - Proprioception training progressions with variable surfaces and perturbations

Criteria for Return to Light Recreational Activity:

1. Full AROM and joint girth at 100% LSI^{16, 17}
2. WB symmetry with squat form to 60°^{16, 17}
3. Stork test at 90% LSI^{16, 17} (Appendix B)
4. Isometric leg press at 60° of knee flexion LSI ≥ 75%^{16, 17} (Appendix C)
5. Isometric quad and HS LSI ≥ 75% at 60° of flexion^{16, 17} (Appendix D-E)
6. Anterior Reach ≤ 4cm difference Vs uninvolved LE^{16, 17} (Appendix F)
7. Single leg hop test LSI ≥ 70%¹⁷ (Appendix G)

Criteria for Progression to Return to Activity Testing:

- Reports confidence with all running and jumping tasks
- Full return to activity should be based on achieving clinical criteria
- If comorbidities create unattainable goals for discharge, discuss this with the treating physician group.

Criteria for Full Return to Recreational/Sport Activity:

General Ortho Patient:

- Patient meets all return to light activity criteria in phase 3.
- Max single leg press LSI ≥ 90%^{9, 10}

Recreational Athlete Sequence (includes above):

- Max Isometric Quad and HS LSI ≥ 90%¹⁰ OKC at 60° of knee flexion.
- Single leg hop test and Crossover hop test²¹ for distance: LSI ≥ 90%¹⁰

Competitive Athlete (includes above):

- Max single leg press LSI ≥ 95%¹⁰
- Max Isometric Quad and HS LSI ≥ 95%¹⁰ OKC at 60° of knee flexion
- Single Leg hop test for distance: LSI ≥ 95%¹⁰
- Side Hop test: LSI ≥ 90%¹⁰ (Appendix G)
- Crossover hop test for distance ≥ 95% LSI^{10, 18} (Appendix I)

Other Literature Review Notes:

- Delayed / decreased outcomes with lateral vs medial partial meniscectomy^{1, 3} potentially due to:
 - Lateral meniscus supports approximately 70% of the load transmission at Tibiofemoral joint¹
 - Lateral meniscus undergoes 2x anteroposterior translation that the medial meniscus does during knee flexion^{3, 11}
 - Higher prevalence of effusion with lateral vs medial meniscectomy with RTS activity¹¹
- Female gender and increased OA before surgery are associated with a slower rate of recovery from arthroscopic partial meniscectomy⁵
- Quad weakness often persists at 6 months and is attributed to neural impairment (activation failure) in maximum concentric and isometric actions. Maximum quad eccentric action did not differ from uninvolved⁷
- Self-reported timeline for return to sport at PLOF:
 - RTS < 30yo 7.7 wks¹²
 - RTS > 30yo 12.7 wks¹²



Abbreviation List:

| | |
|--|--|
| AAROM: Active assisted range of motion | MCL: Medial collateral ligament |
| ABD: Abduction | MD: Medical doctor |
| AD: Assistive device | NWB: Non weight bearing |
| ADL: Activity of daily Living | OKC: Open kinetic chain |
| AROM: Active range of motion | PA: Physician assistant |
| BPTB: Bone patellar tendon bone | PCL: Posterior cruciate ligament |
| BW: Body Weight | PE: Pulmonary embolism |
| CKC: Closed kinetic chain | PLC: Posterior lateral corner |
| DVT: Deep vein thrombosis | PROM: Passive range of motion |
| ER: External rotation | ROM: Range of motion |
| EXT: Extension | RP: Resting position |
| FWB: Full weight bearing | RROM: Resisted range of motion |
| GHJ: Gleno-humeral joint | RTS: Return to sport |
| HEP: Home exercise program | SLR: Straight leg raise |
| HS: Hamstring | UE: Upper extremity |
| IR: Internal rotation | TKE: Terminal knee extension |
| LCL: Lateral collateral ligament | WB: Weight bearing |
| LE: Lower extremity | WBAT: Weight bearing as tolerated |
| MA: Medical assistant | #: Pounds |
| LSI: Limb Symmetry Index = (Average score of the involved leg divided by the score of the uninvolved leg for a specific test) | ≠: Absent/Without |
| | ≈: Approximately |
| | ≤: Less than or equal to |
| | ≥: Greater than or equal to |

Return to Activity Test Descriptions:

Stork Balance Test¹⁹: (Appendix B for diagram)

- Hands on hips. NWB foot: medial distal femur or medial proximal tibia.
- Timer starts when the patient lifts heel of the stance foot off the ground.
- Timer stops if/when the patient removes hands from hips, NWB foot from medial stance leg or the heel comes in contact with the ground.

Anterior Reach Test^{16, 17}: (Appendix F for diagram)

- Stand on one leg and slide a tissue box forward with the toes of the other foot by pushing on the side of the box. Goals is to push the box as far as possible and return back to the starting upright position.
- Once contact is lost between the toes and the box the slide is over.
- Perform 6 warm up attempts per leg to diminish learning effect.
- Failed attempt = the sliding foot touches down on the floor or on top of the slide box before returning back to the starting position. Cannot kick or flick box forwards.
- Distance is measured from toe of standing foot to back edge of the box. Take the best of 3 attempts for each leg.

Single Leg Hop Test for Distance²⁰: (See Appendix D for diagram)

- Measure patient's standing height in cm for pass/fail.
- Hands clasped behind the back to prevent arm swing momentum.
 - Arms can release for landing assistance after leaving ground.
- 4 progressive warm up jumps ≈ 25%, 50%, 75% and 100% intensity.
- Patient must "stick" the landing ≠ significant knee valgus.
- Use the best of 3 maximum effort jump tests.
- Distance is measured from toe of start line to shortest distanced heel.

Single Leg Timed Side Hop Test²¹: (See Appendix E for diagram)

- Set up: 2 parallel lines on floor, with outer edges of lines 40cm apart.
- Start position: standing on single test leg with hands behind the back.
- Action: Patient hops from outside of one line to outside of the other.
- Record the total number of completed foot strikes in 30 seconds.
 - Completed foot strikes = foot lands completely outside the line, without touching the line, while maintaining hand position.

Crossover Hop Test¹⁸: (See Appendix F for diagram)

- Patient starts on one leg with center line just lateral to stance leg.
- Patient is instructed to maximally hop forwards 3 times on the same. stance leg, alternately crossing a ≈15cm wide line.
- Distance is measured from toe of start line to heel of 3rd landed hop.



Quick Reference Activity Timeline:

| Activity | Activity Progression Restrictions |
|--------------------------------------|--|
| Weight Bearing / Gait | <ul style="list-style-type: none"> • Immediately WBAT, wean from crutches without limp as able |
| Knee ROM | <ul style="list-style-type: none"> • 0-90° Day 0-7 then progress flexion as tolerated |
| CKC Squats | <ul style="list-style-type: none"> • As tolerated controlling for effusion, pain and inflammation |
| OKC RROM | <ul style="list-style-type: none"> • As tolerated controlling for effusion, pain and inflammation |
| Plyometrics | <ul style="list-style-type: none"> • ≈ Week 5 With leg press LSI ≥ 75%, OK to begin double leg to single leg with good valgus control • Monitor process to avoid increased swelling/pain |
| Running | <ul style="list-style-type: none"> • Minimum requirement to see leg press LSI ≥ 75% • It is preferable to meet all of the return to light recreational activity criteria on page 3 before running |
| Return to Sport Cleared by MD | <ul style="list-style-type: none"> • Having met the return to activity testing criteria related to level of desired activity intensity on page 3 • Typical return to activity timelines vary ≈ 6-16 weeks^{1, 9, 11, 12} |



Appendix A:
KOOS-pain/KOOS-sport

KOOS-Pain & KOOS-Sport Knee Surveys

Scoring KOOS Tests:
Items are scored on a 0-4 scale. Compare scores from the time of surgery to the time of return to activity to determine if Minimal Clinically Important Difference (MCID) that shows significant positive trend of RTS has been met.

Scoring KOOS-Pain:
The MCID is 9.7 points improvement for KOOS-pain¹

Scoring KOOS-Sport:
The MCID is 14.7 points improvement for KOOS-sport¹

Today's date: ____/____/____ Date of birth: ____/____/____

Name: _____

INSTRUCTIONS: This survey asks for your view about your knee. Answer every question by ticking the appropriate box, only one box for each question. If you are unsure about how to answer a question, please give the best answer you can.

| PAIN: | | | | | |
|---|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| | Never | Monthly | Weekly | Daily | Always |
| 1. How often do you experience pain? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| What amount of knee pain have you experienced the <u>last week</u> during the following activities? | | | | | |
| | None | Mild | Moderate | Severe | Extreme |
| 2. Twisting/pivoting on your knee. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Straightening knee fully. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Bending knee fully. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Walking on flat surface. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Going up or down stairs. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. At night while in bed. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 8. Sitting or lying. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 9. Standing upright. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | | | Total Score 1-9: |

| SPORT: | | | | | |
|---|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| The following questions concern your physical function when being active on a higher level. The questions should be answered thinking of what degree of difficulty you have experienced during the last week due to your knee. | | | | | |
| | None | Mild | Moderate | Severe | Extreme |
| 1. Squatting. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Running. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Jumping. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Twisting/Pivoting on your knee. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Kneeling. | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| | | | | | Total Score 1-5: |



Appendix B: The Stork Test




Maintain hands on hips

Maintain Heel OFF the Ground

Maintain Foot on Medial Leg

Appendix C: Isometric Single Leg Leg Press




60° of Flexion

Dynamometer

- Adjust foot and leg press position so that the knee is in 60 degrees of knee flexion when there is no slack in the dynamometer attachment.
- Perform maximal effort isometric tests per leg.

Appendix D: Isometric Single Leg Quadriceps Test




60° of Flexion

Dynamometer

- Adjust seat position and dynamometer length so that there is no slack in the dynamometer attachment when the knee is in 60° knee flexion.
- Perform max effort isometric tests per leg.

Appendix E: Isometric Single Leg Hamstring







60° of Flexio

Dynamometer

- Adjust seat position and dynamometer length so that there is no slack in the dynamometer attachment when the knee is in 60° knee flexion.
- Perform maximal effort isometric tests per leg.





Meniscus References:

1. Agarwalla, A. et al. Predictive factors and duration to return to sport after isolated meniscectomy. *The Orthopaedic Journal of Sports Medicine*. 2019; 7 (4).
2. Brigham and Women's Hospital, Inc. Department of Rehabilitation Services. Meniscal Repair. 2010.
<https://www.brighamandwomens.org/assets/bwh/patients-and-families/pdfs/knee--partial-meniscectomy-or-debridement.pdf>
3. Brindle, T. et al. The meniscus: Review of Basic Principles with Application to Surgery and Rehabilitation. *Journal of Athletic Training*. 2001; 36(2): 160-169.
4. Ericsson, Y. et al. Muscle Strength, Functional Performance, and Self-Reported Outcomes Four Years After Arthroscopic Partial Meniscectomy in Middle-Aged Patients. *Arthritis and Rheumatism*. December 15 2006; 55(6): 946-952.
5. Fabricant, P. et al. Predictors of Short-term Recover Differ from Those of Long-term Outcomes After Arthroscopic Partial Meniscectomy. *Arthroscopy*. July 2008; 24(7): 769-778.
6. Ford, K et al. Landing Adaptations Following Isolated Lateral Meniscectomy in Athletes. *Knee Surg Sports Traumatol Arthrosc*. Oct 2011; 19(10): 1716-1721.
7. Glatthorn, J. et al. Neuromuscular Function after Arthroscopic Partial Meniscectomy. *Clin Orthop Relat Res*. 2010; 468: 1336-1343.
8. Goodyear-Smith, F. et al. Rehabilitation After Arthroscopic Meniscectomy: A Critical Review of the Clinical Trials. *International Orthopaedics*. 2001; 24: 350-353.
9. Gunderson Health System. Partial Meniscectomy / Chondroplasty Rehabilitation Program. July 2008.
<https://www.gundersenhealth.org/app/files/public/1474/Sports-Medicine-Protocol-Partial-Meniscectomy-Chondroplasty.pdf>
10. Koutras, G. et al. A Randomized Trial of Isokinetic Versus Isotonic Rehabilitation Program After Arthroscopic Meniscectomy. *The International Journal of Sports Physical Therapy*. Feb 2012; 7(1); 31-38.
11. Smoak, J. et al. An Up-to-Date Review of the Meniscus Literature: A Systematic Summary of Systematic Reviews and Meta-analyses *Orthop J Sports Med*. 2020 Sep 9; 8(9): 2325967120950306. doi: 10.1177/2325967120950306. eCollection 2020 Sep.
12. Kim, S. et al. Return to sport after arthroscopic meniscectomy in stable knees. *BMC Sports Science, Medicine and Rehabilitation*. 2013; 5(23).
13. Zhang, X. et al. Effects of Strength and Neuromuscular Training on Functional Performance in Athletes After Partial Medial Meniscectomy. *Journal of Exercise Rehabilitation*. 2017;13(1): 110-116.
14. Wright, R et al. Anterior Cruciate Ligament Reconstruction Rehabilitation: MOON Guidelines. *Sports Health*. 2015 May;7(3):239-43. doi: 10.1177/1941738113517855.
15. Arden, C.L et al. 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of pediatric anterior cruciate ligament injuries. *Knee Surg Sports Traumatol Arthrosc*. 2018 Apr;26(4):989-1010. doi: 10.1007/s00167-018-4865-y. Epub 2018 Feb 17.
16. Gunderson Health System. Meniscus Repair Rehabilitation Program. November 2019.
<https://www.gundersenhealth.org/app/files/public/1473/Sports-Medicine-Protocol-Meniscus-Repair.pdf>



17. Harput, G. et al. Postoperative rehabilitation and outcomes following arthroscopic isolated meniscus repairs: A systematic review. *Physical Therapy in Sport*. 2020 Sep;45:76-85. doi: 10.1016/j.ptsp.2020.06.011. Epub 2020 Jul 13
18. Munro, A et al. Between-Session Reliability of Four Hop Tests and the Agility T-Test. *Journal of strength and conditioning research*. May 2011; 25, (5): 1470-1477.
19. Schell, J et al. *Physical Fitness Assessment in Exercise and Sports Science*. 1994 2nd Ed, Leelar Biomedisience Services, Matraville, NSW. p. 327
20. Davies G.J et al. Individualizing the Return to Sports After Anterior Cruciate Ligament Reconstruction. *Operative Techniques in Orthopaedics*. 2017 Mar 27:1 70-78.
21. Gustavsson, A. et al. A test battery for evaluating hop performance in patients with an ACL injury and patients who have undergone ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2006; 14: 778-788. DOI 10.1007/s00167-006-0045-6.

ValleyOrtho

 VALLEY VIEW