

ValleyOrtho Rehabilitation Playbook Series

Physician: Dr. Chris George
Office Phone: 970-384-7140

Physician Assistant: Rachel Mazza
Office Fax: 970-384-8133

MA: Michelle Gorton

ATC: Jackie Brey

Surgical Procedure: Total Knee Replacement

The intent of this information is to inform the treating clinician on the evidence-based considerations to be used as a guideline regarding the surgery noted above. This is not a substitute for appropriate clinical decision making, but a supplement to that effect. If at any time a clinician feels uncertain about a given phase discrepancy or patient presentation they are strongly encouraged to discuss this with the referring physician and his/her team before alarming the patient. The goal of this rehabilitation guideline IS NOT to be used to motivate patients through fear and discouragement if they are not attaining goals in the described timeframes but to increase physician-therapist communication around established principles. It is the responsibility of the therapist to read the operative report before providing care to the patient to improve treatment communication.

Therapeutic Activity and Phase Progression Disclaimer: Progression to the next phase should be strongly based on meeting clinical criteria and goals of the previous phase (not solely based on the post-operative timeframes) as appropriate and in collaboration with the referring surgeon. Exercise prescription should be clinically directed by pain and performance absent of detrimental compensation with respect to proper biomechanics of the spine, hip, knee and ankle.

Communication Recommendations from Therapist to Surgical Team:

When a treating therapist feels the need to reach out to Dr. George, or a member of his team, at any point for any reason, they are strongly encouraged to do so. All concerns are not explicitly written and clinical judgement is paramount. Below is a handful of reasons and suggested methods of contact to promote communication:

Urgent Red Flag Communication: the patient is in clinic and an action is required as directed by referring staff office

- Uncontrollable and unremitting pain
- Signs of infection at incision or treated limb
- Severe palpation tenderness, swelling, tachycardia (UE or LE DVT)
- Labored breathing (PE)
- Drastic improvement or decline in ROM (failed component)
- Severe pain with walking or ROM
- After a fall/trauma, or near fall/trauma, resulting in a clinical change

Preferred Contact Method: Immediate phone call to speak with MA or ATC until answer.

Administrative Needs

- Rehabilitation Prescription needed or prescription change requests
- Appointment needed with the physician office, or medication refill

Preferred Contact Method: Phone call to MA/ATC

Other Patient Concerns During Clinic Hours M-TH 9-5pm F 9-3pm

- Abnormal pain, comorbidities or complications that may prevent attainment of established discharge criteria
- Patient is noncompliant with rehabilitation process
- Excessive muscle guarding/motion phobia after 1-2 outpatient visits
- Adverse work or home practices negatively impacting recovery
- Patient expresses discontent or concerns with the current POC established by PT and/or by MD/PA

Preferred Contact Method: Phone call to MD &/or PA

Updates before checkup visits with MD/PA

During Clinic Hours M-TH 9am-5pm F 9am-3pm

- Information regarding adherence/participation in rehabilitation process
- Comments on progress and trending nature of the patient's rehab course

Preferred Contact Method: Phone call to MD &/or PA. Or Fax update



Phase 1: Post-Op Healing and Knee Extension (Wks 1 & 2)

Goals:

- Reduce swelling and pain
- Patient adheres to pain management strategy from physician
- Independent transfers and gait (with or without AD)
- Knee AROM 5°-90°, Knee extension PROM to 0°
- Patient able to maintain a quad set for 6 seconds at end range ext
- Ensure appropriate compression for 2 weeks (if TED hose are creating a tourniquet adjust to alternative type: Tubigrip/Knee highs or Ace Wrap)

Precautions:

- No ambulation without AD if gait deviations are present
- Patient should avoid static knee flexion positions for >15 minutes at a time until full extension is well established.
- Avoid too little or too much use of pain medications
- Avoid activity that results in increased swelling lasting > 24 hours
- Incision should not be submerged until full closure without scabbing

Phase 1 Therapeutic Activities:

- Aggressive edema control (compression, ice, elevation, soft tissue work)
- Quad, glute, hamstring and calf activation and isometric exercises
- NMES to Quads with SAQ/QS if having difficulty with contraction
- A/AA/PROM/joint mobilization into knee extension
- AAROM knee flexion
- Educate ADL strategies for tub/shower/car/ etc.
- Gait training on level surface with least restrictive AD for normal pattern
- Nonreciprocal stair training

Criteria for Progression to Phase 2:

- Attain phase 1 goals

Phase 2: ROM and Early Strengthening (Weeks 3 & 4)

Goals:

- Normalized gait to pre-injury AD status
- Independent with HEP and ADLs
- Knee AROM <5°-100° or more
- Knee extension PROM to 0°

Precautions:

- Phase 1 precautions included
- Avoid pain increase greater than 5/10 with exercise or functional activity to avoid muscle inhibition

Phase 2 Therapeutic Activities:

- Cryotherapy and edema control PRN
- Initiate stationary bike and core exercises
- A/PROM stretching to hamstrings, quads and calf
- Manual work into knee extension unless AROM extension is at 0°
- Quad tendon, patellar tendon and patella superior + inferior mobilizations
- Scar mobilizations
- CKC TKE exercises
- Balance/proprioception training progressing from double to single leg
- CKC partial body weight through available knee ROM keeping below 5/10 pain

Criteria for Progression to Phase 3:

- Attain phase 2 goals



Phase 3: Intermediate Strengthening (Weeks 5 & 6)

Goals:

- Knee AROM 0°-110° or more
- SLR without quad lag
- Hip strength equal to non-operative side
- Reciprocal with stairs

Precautions:

- F/U with MD if plateau in ROM, constant pain and/or maintained edema
- No twisting /pivoting until >12 weeks
- Caution with moderate impact activities until > 12 weeks

Phase 3 Therapeutic Activities:

- 4 way resisted SLR (OKC and CKC) for hip strengthening
- Manual work into knee extension unless AROM extension is at 0°
- Manual work into knee Flexion as necessary for phase goal
- Single leg balance progressions
- CKC full body weight through available knee ROM

Criteria for Progression to Phase 4:

- Attain phase 3 goals

Phase 4: Advanced Strength and Final HEP (Wks 7 to D/C)

Goals:

- Return to full pre injury function excluding high impact pivot/twisting activities
- Meet discharge criteria

Precautions:

- Same as phase 3
- Pain after activity lasting >24 hours is not encouraged

Phase 4 Therapeutic Activities:

- Increase resistance, reps and sets for all exercises
- Include specific return to job or sport specific training
- Transition to appropriate HEP
- Ensure patient is independent getting from floor to standing

Minimum Low Activity Patient D/C Criteria:

- Return to all normal ADLs and light recreational activities without pain
- Able to get on to and off of the floor with minimal use of hands
- Normalized gait pattern
- Reciprocal pattern with ascending and descending stairs
- Knee AROM 0°-120° or more
- Patient is consistent and independent with HEP
- Single leg stance of 5 seconds with eyes open and TUG of < 13.5 seconds

High Activity Patient D/C Criteria (includes above):

- Limb symmetry of 90% or greater in strength and ROM
- Single leg stance time with eyes closed based on age:
(50-59yo = 25 seconds) (60-69yo = 15 seconds) (70-79yo = 6 seconds)



Abbreviation List:

AAROM: Active assisted range of motion

AD: Assistive device

ADL: Activity of daily Living

AROM: Active range of motion

BW: Body Weight

CKC: Closed kinetic chain

DVT: Deep vein thrombosis

D/C: Discharge

ER: External rotation

EXT: Extension

FWB: Full weight bearing

F/U: Follow up

HEP: Home exercise program

IR: Internal rotation

LE: Lower extremity

MA: Medical assistant

MD: Medical doctor

NWB: Non weight bearing

OKC: Open kinetic chain

PA: Physician assistant

PE: Pulmonary embolism

PROM: Passive range of motion

ROM: Range of motion

RP: Resting position

RROM: Resisted range of motion

SLR: Straight leg raise

UE: Upper extremity

WB: Weight bearing

YO: Years old

≈: Approximate

#: Pounds

References

1. Artz Net et al. Effectiveness of physiotherapy exercise following total knee replacement: systematic review and meta-analysis. *BMC Musculoskelet Disord.* 2015 Feb 7;16:15.
2. Bandholm T et al. Rehabilitation strategies for optimisation of functional recovery after major joint replacement. *J Exp Orthop.* 2018 Oct 11;5(1):44.
3. Brigham and Women's Hospital Department of Rehabilitation Services, Total Knee Arthroplasty protocol. <https://www.brighamandwomens.org/patients-and-families/rehabilitation-services/physical-therapy-standards>.
4. Cahill, JB. Postsurgical Rehabilitation Guidelines for the Orthopedic Clinician. June 2006.
5. Cincinnati Sports medicine and orthopaedic center, Total knee replacement: rehabilitation protocol*. https://cincinnati-sportsmed.com/wp-content/uploads/2011/03/Total_Knee_Replacement.pdf
6. Florez-García M et al. Efficacy and safety of home-based exercises versus individualized supervised outpatient physical therapy programs after total knee arthroplasty: a systematic review and meta-analysis. *Knee Surg Sports Traumatol Arthrosc.* 2017 Nov;25(11):3340-3353.
7. George, C. M.D. Expert opinion and consultation.
8. Husted RS et al. Rapid knee extensions to increase quadriceps muscle activity in patients with total knee arthroplasty: a randomized cross-over study. *Int J Sports Phys Ther.* 2017 Feb;12(1):105-116.
9. Schache MB et al. Does the addition of hip strengthening exercises improve outcomes following total knee arthroplasty? A study protocol for a randomized trial. *BMC Musculoskelet Disord.* 2016 Jun 13;17:259.
10. The Stone Clinic, Total Knee Replacement Post-Operative Rehabilitation Protocol. <https://www.stoneclinic.com/total-knee-replacement-rehab-protocol>.



11. Lahey Hospital and Medical Center, Rehabilitation Protocol: Total Knee Arthroplasty. <https://www.lahey.org/RehabilitationProtocols/>.
12. Anderson, M. et al. Fall risk and utilization of balance training for adults with symptomatic knee osteoarthritis: Secondary analysis from a randomized control trial. *Journal of Geriatric Physical Therapy*. 2018.
13. Bohannon et al. Decrease in timed balance test scores with aging. *Phys Ther*. 1984; 64:1067-1070.
14. Snell, D et al. Rehabilitation after total joint replacement: a scoping study. *Disability and Rehabilitation*, 2018 Jul;40(14):1718-1731.

ValleyOrtho

